### For Research Use Only

# HNRNPH1 Monoclonal antibody

Catalog Number:67375-1-lg 2 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

BC001348

Size:

GeneID (NCBI):

1000 ug/ml

3187

Source:

UNIPROT ID:

Mouse

P31943

Isotype:

GenBank Accession Number:

BC001348

UNCBI):

1000 ug/ml

3187

Full Name:

IgG1 heterogeneous nuclear ribonucleoprotein H1 (H)

AG6603 Calculated MW: 49 kDa

Observed MW: 49-50 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

Species Specificity: human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, MCF-7 cells, LNCaP cells, U2OS cells, HSC-T6 cells, 4T1 cells, Jurkat cells,

**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:500-1:2000

IF/ICC 1:50-1:500

CloneNo.:

1A6C2

NIH/3T3 cells

IHC: human breast cancer tissue,

IF/ICC: HepG2 cells,

## **Background Information**

The heterogeneous nuclear ribonucleoprotein (hnRNP) complexes which provide the substrate for the processing events that pre-mRNAs undergo before becoming functional, translatable mRNAs in the cytoplasm. HNRNPH1 is one component of hnRNP complex and mediates pre-mRNA alternative splicing regulation [PMID:11003644]. Together with CUGBP1, it inhibits IR pre-mRNA exon 11 inclusion in myoblast [PMID: 16946708].

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Shiqing Wang	39221900	J Cell Physiol	WB
Tamara Vital	36793594	Front Oncol	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 67375-1-1g (HNRNPH1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67375-1-1g (HNRNPH1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67375-1-lg (HNRNPH1 antibody), at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).