For Research Use Only

TSG101 Monoclonal antibody

Catalog Number:67381-1-lg 47 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 67381-1-lg
 BC002487

 Concentration:
 GeneID (NCBI):

 1000 μ g/ml
 7251

Source: UNIPROT ID:
Mouse Q99816
Isotype: Full Name:
IgG1 tumor suscept

Immunogen Catalog Number: Calculated MW:

AG28569 44 kDa

Observed MW: 43-46 kDa

tumor susceptibility gene 101

Purification Method:

2B7G8

Protein G purification CloneNo.:

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000 IF-P 1:200-1:800

IF/ICC 1:1000-1:4000

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications:

WB, IF

Species Specificity: human, mouse, rat Cited Species:

human, mouse, rat, rabbit, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: LNCaP cells, HSC-T6 cells, human urine exosomes tissue, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, C6 cells, NIH/3T3 cells

 $\label{lhc:human} \textbf{IHC:} human \, lung \, cancer \, tissue, \, human \, colon \, cancer$

issue

IF-P: human colon cancer tissue,

IF/ICC: HeLa cells, human colon cancer tissue

Background Information

TSG101(Tumor susceptibility gene 101 protein) is essential for endosomal sorting, membrane receptor degradation and the final stages of cytokinesis. It plays a crucial role for cell proliferation and cell survival. TSG101 has been identified as a candidate tumor suppressor gene and belongs to the ubiquitin-conjugating enzyme family. TSG101 is a marker for exosome. This protein has 2 isoforms produced by alternative splicing with the molecular mass of 44 and 32 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhaoyi Liang	36210461	Cell Mol Biol Lett	WB
Hang Yuan	36276139	Front Oncol	WB
Xiaolu Huang	36426920	J Clin Lab Anal	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67381-1-lg (TSG101 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control




Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using TSG101 antibody (67381-1-lg, Clone: 2B7G8) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67381-1-Ig (TSG101 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



 $Immun of luorescent analysis of (-20^{\circ}C \ Ethanol) fixed \ HeLa cells using 67381-1-lg (TSG101 \ antibody), at dilution of 1:2000 and Coralite @488-Conjugated Goat Anti-Mouse IgG(H+L).$



human urine exosomes tissue were subjected to SDS PAGE followed by western blot with 67381-1-Ig (TSG101 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.