For Research Use Only

AXIN2 Monoclonal antibody

Catalog Number:67458-1-lg




Basic Information	Catalog Number: 67458-1-lg	GenBank Accession Number: BC006295	Purification Method: Protein G purification
	Concentration: 1100 ug/ml	GenelD (NCBI): 8313	CloneNo.: 1A11A4
	Source: Mouse	UNIPROT ID: Q9Y2T1	Recommended Dilutions: WB 1:1000-1:9600
	lsotype: lgG1	Full Name: axin 2	
	Immunogen Catalog Number: AG29291	Calculated MW: 843 aa, 94 kDa	
		Observed MW: 100 kDa	
Applications	Tested Applications:		
	WB, ELISAWB : MDASpecies Specificity:12 cellshuman, rat12 cells		-MB-231 cells, HepG2 cells, Jurkat cells, PC
	Axis inhibition protein2 (AXIN2), also known as Coductin or Axil, is a multidomain scaffold protein that negatively regulate Wnt signaling. AXIN2 can directly interact with beta-catenin and GSK3B. AXIN2 has a number of phosphorylation sites, and also can undergo poly(ADP-ribosy)lation by tankyrase TNKS and TNKS2. Poly(ADP- ribosy)lated AXIN2 then get unbiquitinated by RNF146 and leads to its degradation and subsequent activation of Wnt signaling. AXIN2 is localized in the cytoplasm and its mutation is involved in colorectal cancer.		
Background Information	regulate Wnt signaling. AXIN2 car phosphorylation sites, and also ca ribosy)lated AXIN2 then get unbic	n directly interact with beta-catenin an n undergo poly(ADP-ribosy)lation by juitinated by RNF146 and leads to its	nd GSK3B. AXIN2 has a number of tankyrase TNKS and TNKS2. Poly(ADP- degradation and subsequent activation of
Background Information	regulate Wnt signaling. AXIN2 car phosphorylation sites, and also ca ribosy)lated AXIN2 then get unbic	n directly interact with beta-catenin and n undergo poly(ADP-ribosy)lation by uitinated by RNF146 and leads to its in the cytoplasm and its mutation is i after shipment.	nd GSK3B. AXIN2 has a number of tankyrase TNKS and TNKS2. Poly(ADP- degradation and subsequent activation of

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67458-1-1g (AXIN2 antibody) at dilution of 1:4800 incubated at room temperature for 1.5 hours.