

For Research Use Only

SF3B3 Monoclonal antibody

Catalog Number: 67469-1-Ig **1 Publications**



Basic Information

Catalog Number: 67469-1-Ig	GenBank Accession Number: BC003146	Purification Method: Protein A purification
Size: 1400 ug/ml	GeneID (NCBI): 23450	CloneNo.: 2H7F5
Source: Mouse	UNIPROT ID: Q15393	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2b	Full Name: splicing factor 3b, subunit 3, 130kDa	IHC 1:1000-1:4000
Immunogen Catalog Number: AG6315	Calculated MW: 136 kDa	IF-P 1:200-1:800
	Observed MW: 130-135 kDa	

Applications

Tested Applications: WB, IHC, IF-P, ELISA	Positive Controls:
Cited Applications: IF	WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, RAW 264.7 cells
Species Specificity: human, mouse, rat	IHC : human breast cancer tissue, human gliomas tissue
Cited Species: mouse	IF-P : mouse heart tissue, rat heart tissue

Note-IHC: suggested antigen retrieval with *TE buffer pH 9.0*; (*) Alternatively, antigen retrieval may be performed with *citrate buffer pH 6.0*

Background Information

Introns are removed from nuclear pre-mRNA in 2-step transesterification reactions. Splicing occurred in a large ribonucleoprotein particle, called the spliceosome. Spliceosomal intermediate complexes form on pre-mRNA in the order E, A, B, and C, with the catalytic reactions occurring in complex C. U2 small nuclear ribonucleoproteins are one of the proteins essential for spliceosome assembly and mRNA splicing. Functional U2 snRNP is composed of a 12S unit and 2 splicing factors, SF3A, which is composed of 3 proteins, and SF3B, which composed of 4 proteins. SF3B3 is one of SF3B, and it's required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence(BPS) in pre-mRNA.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Cui	38188517	iScience	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67469-1-Ig (SF3B3 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67469-1-Ig (SF3B3 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat heart tissue using SF3B3 antibody (67469-1-Ig, Clone: 2H7F5) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse heart tissue using SF3B3 antibody (67469-1-Ig, Clone: 2H7F5) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).