For Research Use Only

PGAM1 Monoclonal antibody

Catalog Number:67470-1-lg Featured Product

2 Publications

BC011678

5223

P18669

GeneID (NCBI):

UNIPROT ID:

Full Name:

Calculated MW:

GenBank Accession Number:

phosphoglycerate mutase 1 (brain)



Basic Information

Catalog Number: 67470-1-lg

Size: 1000 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG9250

254 aa, 29 kDa Observed MW:

29 kDa

Positive Controls:

WB: U2OS cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, A549 cells, LNCaP cells, K-562 cells

Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:200-1:1000

IF/ICC 1:400-1:1600

CloneNo.:

2H2A9

IHC: human liver cancer tissue.

IF/ICC: HepG2 cells,

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity: Human, Mouse, Rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PGAM1(phosphoglycerate mutase 1) is also named as PGAMA, PGAM-B and belongs to the phosphoglycerate mutase family. Phosphoglycerate mutase is widely distributed in mammalian tissues where it catalyzes the reversible reaction of 3-phosphoglycerate (3-PGA) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The homodimer PGAM1 is expressed mainly in liver, kidney, brain and overexpressed in a variety of human cancers, including breast carcinoma, colorectal cancer, lung cancer, prostate cancer, oral squamous cell carcinoma, esophageal squamous cell carcinomas and also associated with certain virus infection. PGAM1 could be developed as a useful diagnostic biomarker, as well as a potential therapeutic target for hepatocellular carcinoma (PMID:20403181). This antibody may also recognize PGAM2 and PGAM4 due to the high homology.

Notable Publications

Author	Pubmed ID	Journal	Application
Wei Zhang	34689761	J Nanobiotechnology	WB,IHC
Xinlu Liu	29386088	Oncol Res	IF

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




U2OS cells were subjected to SDS PAGE followed by western blot with 67470-1-1g (PGAM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67470-1-Ig (PGAM1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67470-1-Ig (PGAM1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PGAM1 antibody (67470-1-lg, Clone: 2H2A9) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).