

For Research Use Only

PGAM1 Monoclonal antibody, PBS Only



Catalog Number: 67470-1-PBS

Featured Product

Basic Information

Catalog Number:

67470-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG9250

GenBank Accession Number:

BC011678

GeneID (NCBI):

5223

UNIPROT ID:

P18669

Full Name:

phosphoglycerate mutase 1 (brain)

Calculated MW:

254 aa, 29 kDa

Observed MW:

29 kDa

Purification Method:

Protein G purification

CloneNo.:

2H2A9

Applications

Tested Applications:

WB, IHC, IF, ELISA

Species Specificity:

Human, Mouse, Rat

Background Information

PGAM1(phosphoglycerate mutase 1) is also named as PGAMA,PGAM-B and belongs to the phosphoglycerate mutase family. Phosphoglycerate mutase is widely distributed in mammalian tissues where it catalyzes the reversible reaction of 3-phosphoglycerate (3-PGA) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The homodimer PGAM1 is expressed mainly in liver, kidney, brain and overexpressed in a variety of human cancers, including breast carcinoma, colorectal cancer, lung cancer, prostate cancer, oral squamous cell carcinoma, esophageal squamous cell carcinomas and also associated with certain virus infection. PGAM1 could be developed as a useful diagnostic biomarker, as well as a potential therapeutic target for hepatocellular carcinoma (PMID:20403181). This antibody may also recognize PGAM2 and PGAM4 due to the high homology.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com




W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



U2OS cells were subjected to SDS PAGE followed by western blot with 67470-1-Ig (PGAM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67470-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PGAM1 antibody (67470-1-Ig, Clone: 2H2A9) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67470-1-PBS in a different storage buffer formulation.