

For Research Use Only

ASC/TMS1 Monoclonal antibody

Catalog Number: 67494-1-Ig **31 Publications**



Basic Information

Catalog Number: 67494-1-Ig	GenBank Accession Number: BC004470	Purification Method: Protein A purification
Concentration: 1000 ug/ml	GeneID (NCBI): 29108	CloneNo.: 2G12E6
Source: Mouse	UNIPROT ID: Q9ULZ3	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: PYD and CARD domain containing	
Immunogen Catalog Number: AG28424	Calculated MW: 25 kDa	
	Observed MW: 22-25 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HL-60 cells, human placenta tissue, THP-1 cells, U-937 cells
Cited Applications: WB, IHC, IF, IP, CoIP	
Species Specificity: human	
Cited Species: human	

Background Information

PYCARD, also known as ASC or TMS1, is a 195 amino acid protein containing an N-terminal Pyrin-like domain (PYD) and an 87 residue C-terminal Caspase-associated recruitment domain (CARD). It promotes caspase-mediated apoptosis which is mediated predominantly through the activation of caspase-9.

Notable Publications

Author	Pubmed ID	Journal	Application
Jiayang Liu	34733914	Ann Transl Med	WB
Yiping Leng	32912629	Biochem Biophys Res Commun	WB
Yingchun Zhao	34663844	Sci Rep	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67494-1-Ig (ASC/TMS1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



human placenta tissue were subjected to SDS PAGE followed by western blot with 67494-1-Ig (ASC/TMS1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.