For Research Use Only

PER2 (isoform 1) Monoclonal antibody

Catalog Number:67513-1-lg 12 Publications



Basic Information

Applications

Catalog Number: 67513-1-lg

Concentration: 1000 ug/ml Source: Mouse

Isotype: lgG1

Immunogen Catalog Number: AG29489

Tested Applications: WB, IF/ICC, ELISA **Cited Applications:**

WB, IHC

Species Specificity:

human **Cited Species:** human, mouse, pig GenBank Accession Number:

NM_022817 GeneID (NCBI): 8864

UNIPROT ID: 015055 Full Name:

period homolog 2 (Drosophila)

Calculated MW: 137 kDa Observed MW:

150~160 kDa

Positive Controls:

WB: HEK-293 cells, LO2 cells, Y79 cells, BxPC-3 cells, K-

Purification Method:

CloneNo.:

2E6C3

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:400-1:1600

562 cells, HL-60 cells, THP-1 cells

IF/ICC: HeLa cells.

Background Information

PER2, also named as KIAA0347, is a component of the circadian clock mechanism which is essential for generating circadian rhythms. Defects in PER2 are a cause of familial advanced sleep-phase syndrome (FASPS) which is characterized by very early sleep onset and offset. There are two isoforms of PER2: isoform 1 has an expected molecular size of about 140 kDa, and 67513-1-Ig detected molecular weight of 150-160 kDa (PMID:16675517), while isoform 2, known as PER2S (a second splicing variant of the human PER2 gene), has a molecular weight of about 45 kDa (PMID:26347774).

Notable Publications

Author	Pubmed ID	Journal	Application
Xuekai Xiong	36093830	J Cell Sci	WB
Zhaoxia Wang	33244267	Cancer Manag Res	WB
Xiyan Tan	34124276	J Immunol Res	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67513-1-1g (PER2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PER2 (isoform 1) antibody (67513-1-lg, Clone: 2E6C3) at dilution of 1:800 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).