

For Research Use Only

HPRT Monoclonal antibody, PBS Only

Catalog Number: 67518-1-PBS

Featured Product



Basic Information

Catalog Number:

67518-1-PBS

Concentration:

1mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG7044

GenBank Accession Number:

BC000578

GeneID (NCBI):

3251

UNIPROT ID:

P00492

Full Name:

hypoxanthine
phosphoribosyltransferase 1

Calculated MW:

25 kDa

Observed MW:

24-28 kDa

Purification Method:

Protein A purification

CloneNo.:

1C3D1

Applications

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

HPRT, also named HPRT1 and HGPRT, plays a central role in the generation of purine nucleotides through the purine salvage pathway. Mutation of HPRT1 is associated with Lesch-Nyhan syndrome (LNS) which is an X-linked inherited neurogenetic disorder of purine metabolism. It has been reported that HPRT1 also plays an important role in HPRT-related gout. 67518-1-Ig antibody detects the protein around 28 kDa in SDS-PAGE. And HPRT1 deficient cell line Sp2/0 was served as a negative control in PTC's results. (PMID:20544509, 28045594, 27379977)

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67518-1-Ig (HPRT antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. HPRT deficient cell line Sp2/0 was served as a negative control. This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 67518-1-Ig (HPRT antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 67518-1-Ig (HPRT antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human endometrial cancer tissue slide using 67518-1-Ig (HPRT antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human endometrial cancer tissue slide using 67518-1-Ig (HPRT antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using HPRT antibody (67518-1-Ig, Clone: 1C3D1) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.



WB result of HPRT antibody (67518-1-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HPRT transfected HEK-293 cells. This data was developed using the same antibody clone with 67518-1-PBS in a different storage buffer formulation.