For Research Use Only

## CCT8 Monoclonal antibody

Catalog Number:67539-1-lg Featured Product 4 Publications






Basic Information	Catalog Number: 67539-1-lg	GenBank Accession Number: BC012584		Purification Method: Protein A purification	
	Size: 1700 µg/ml	GeneID (NCBI): 10694		CloneNo.: 1A7C3	
	Source: Mouse	UNIPROT ID: P50990		Recommended Dilutions: WB 1:5000-1:50000	
	Isotype: IgG2a Immunogen Catalog Number: AG30082	Full Name: chaperonin conta 8 (theta)	chaperonin containing TCP1, subunit IF/ICC 1:750-1:3000		
		Calculated MW: 548 aa, 60 kDa Observed MW: 60 kDa			
Applications	Tested Applications:		Positive Controls:		
	Cited Applications: cells, Ju WB IHC IP			ells, NIH/3T3 cells, A549 cells, HeLa Ills, K-562 cells, HSC-T6 cells	
			IHC : human li	IHC : human liver cancer tissue,	
	Human, mouse, rat IF/ICC : MCF-7 cells,			' cells,	
	Cited Species: human				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	chaperonin family. It is a molec BBS/CCT complex, CCT8 may pl	ular chaperone; assists t lay a role in the assemb	he folding of protein ly of BBSome, a com	nd NY-REN-15, belongs to the TCP-1 is upon ATP hydrolysis. As part of the plex involved in ciliogenesis regular liferation and invasion of tumors.	
Notable Publications	Author	Pubmed ID Jo	ournal	Application	
	Xinyu Ma	35366418 C	ell	WB	
	Ruijuan Du	39259093 M	edicine (Baltimore)	IHC	
	Zhiqiang Wu	37928427 J	Cancer	WB,IHC,IP	
Storage	Storage: Store at -20°C. Stable for one ye Storage Buffer:	ar after shipment.			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data


temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 67539-1-Ig (CCT8 antibody) at dilution of 1:20000 incubated at room




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67539-1-1g (CCT8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67539-1-1g (CCT8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CCT8 antibody (67539-1-lg, Clone: 1A7C3) at dilution of 1:1500 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



WB result of CCT8 antibody (67539-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCT8 transfected HEK-293 cells.