For Research Use Only

## EEF2 Monoclonal antibody

Catalog Number:67550-1-lg




Basic Information	Catalog Number: GenBank Accession Number: 57550-1-lg BC 126259		r: Purification Method: Protein G purification	
	Concentration: 1000 µg/ml	GeneID (NCBI): 1938	CloneNo.: 1E4D5	
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG13859	UNIPROT ID: P13639	Recommended Dilutions: WB 1:5000-1:50000	
		Full Name: eukaryotic translation elor factor 2	IHC 1:500-1:2000 Igation IF/ICC 1:400-1:1600	
		Calculated MW: 858 aa, 95 kDa		
		Observed MW: 100 kDa		
Applications	Tested Applications: WB, IHC, IF/ICC, ELISA		Positive Controls: WB : HeLa cells, HSC-T6 cells, LoVo cells, NIH/3T3 cells, HEK-293 cells, HepG2 cells, Jurkat cells, 4T1 cells, MCF-7 cells, pig brain tissue	
	Species Specificity: human, mouse, rat, pig			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		: human breast cancer tissue, human colon cance ue	
			CC : HepG2 cells,	
Background Information	EEF2 (Eukaryotic Elongation Factor 2), also known as EF-2 and EF2, is a crucial protein involved in the process of protein synthesis in eukaryotic cells. It plays a pivotal role in the elongation phase of translation, where it facilitates the movement of the ribosome along the mRNA strand. This movement is essential for the addition of amino acids to the growing polypeptide chain.			
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage			


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67550-1-lg (EEF2 antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67550-1-1g (EEF2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 67550-1-lg (EEF2 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using EEF2 antibody (67550-1-Ig, Clone: 1E4D5) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).