For Research Use Only

RPS14 Monoclonal antibody

Catalog Number:67566-1-lg Featured Product



Basic Information

Catalog Number: 67566-1-Ig

Size: 2000 ug/ml

Source: Mouse

Isotype:

lgG2b

GenBank Accession Number: BC003401 GeneID (NCBI): 6208

UNIPROT ID: P62263 Full Name: ribosomal protein S14

Immunogen Catalog Number: Calculated MW:

AG30032 16 kDa

Observed MW: 16 kDa Purification Method:

Protein A purification

CloneNo.: 3D5H5

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:400-1:1600

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A549 cells, HeLa cells, HepG2 cells, HuH-7 cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, 4T1

cells

IHC: human urothelial carcinoma tissue,

IF/ICC: HepG2 cells,

Background Information

The mammalian ribosome consists of 4 RNA species and approximately 80 different proteins. The ribosomal proteins are encoded by complex gene families that include at least 1 active intron-containing gene and multiple processed pseudogenes[PMID 3785212]. The RPS14 gene, encoding a ribosomal protein that belongs to the 40S ribosome, and mapping to the commonly deleted region (CDR) of the 5q-syndrome, shows haploinsufficiency in the CD34+ cells of patients with this disorder[PMID:22430631].

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67566-1-lg (RPS14 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control




Immunohistochemical analysis of paraffinembedded human urothelial carcinoma tissue slide using 67566-1-Ig (RPS14 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 67566-1-1g (RPS14 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of RPS14 antibody (67566-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RPS14 transfected HeLa cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RPS14 antibody (67566-1-lg, Clone: 3D5H5) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).