

For Research Use Only

STAT4 Monoclonal antibody

Catalog Number: 67568-2-Ig **2 Publications**



Basic Information

Catalog Number: 67568-2-Ig	GenBank Accession Number: BC031212	Purification Method: Protein G purification
Size: 2000 ug/ml	GeneID (NCBI): 6775	CloneNo.: 4A8C9
Source: Mouse	UNIPROT ID: Q14765	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000 IF/ICC 1:50-1:500
Isotype: IgG1	Full Name: signal transducer and activator of transcription 4	
Immunogen Catalog Number: AG19545	Calculated MW: 748 aa, 86 kDa	
	Observed MW: 86 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : HeLa cells, 4T1 cells, HEK-293 cells, HepG2 cells, Jurkat cells, MOLT-4 cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells
Cited Applications: WB, IHC, IF	IHC : human breast cancer tissue, human cervical cancer tissue
Species Specificity: human, mouse, rat	IF/ICC : HepG2 cells,
Cited Species: mouse, rat	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The JAK/STAT pathway is an extensive signaling pathway downstream of cytokine receptors. STATs are cytosolic proteins with a common structure consisting of an N-terminal oligomerization domain, which favors formation of STAT dimers, followed by a DNA-binding domain and a C-terminal SRC homology-2 (SH2) domain, which is involved in association between STATs and receptors[PMID:22383755]. Signal Transducer and Activator of Transcription 4 (STAT4) is a transcription factor that is activated by IL-12 signaling and promotes Th1-cell differentiation and IFN- γ production [PMID:21998209].

Notable Publications

Author	Pubmed ID	Journal	Application
Jiannan Zhou	38574973	Free Radic Biol Med	IF
Xi-Yun Liu	37875707	Mol Neurobiol	WB,IHC


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67568-2-Ig (STAT4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67568-2-Ig (STAT4 antibody), at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67568-2-Ig (STAT4 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67568-2-Ig (STAT4 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug STAT4 Monoclonal antibody (67568-2-Ig, Clone:4A8C9) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1) (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).