For Research Use Only

## CDK1 Monoclonal antibody, PBS Only

Catalog Number:67575-1-PBS




Basic Information	Catalog Number: 67575-1-PBS	GenBank Accession Number: BC014563	Purification Method: Protein A purification
	Size:	GeneID (NCBI):	CloneNo.:
	1mg/ml	983	2A11E4
	Source: Mouse	UNIPROT ID: P06493	
	lsotype: IgG2a	Full Name: cell division cycle 2, G1 to S and G2 to M	
	Immunogen Catalog Number: AG29018		
		Calculated MW: 25 kDa	
		Observed MW: 30-34 kDa	
Applications	Tested Applications: WB, FC (Intra), ELISA		
	Species Specificity: human, mouse, rat		
Background Information	CDK1, also named as CDC2, belongs to the protein kinase superfamily, CMGC Ser/Thr protein kinase family and CDC2/CDKX subfamily. CDK1 plays a key role in the control of the eukaryotic cell cycle. It is required in higher cells for entry into S-phase and mitosis. CDK1 is a catalytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), which is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. It is a component of the kinase complex that phosphorylates the repetitive C-terminus of RNA polymerase II. Mitotic cyclins stably associate with CDK1 and function as regulatory subunits. CDK1 has 2 isoforms produced by alternative splicing with the molecular mass of 34 kDa and 27 kDa.		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67575-1-Ig (CDK1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67575-1-PBS in a different storage buffer formulation. 1x10^6 HeLa cells were intracellularly stained with 0.25 ug CDK1 Monoclonal antibody (67575-1-1g, Clone:2A11E4) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG2a isotype control Mouse McAb (66360-2-Ig, Clone: 11A1B2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67575-1-PBS in a different storage buffer