

For Research Use Only

SLC25A17 Monoclonal antibody

Catalog Number: 67635-1-Ig



Basic Information

Catalog Number:

67635-1-Ig

Size:

1000 μ g/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG24456

GenBank Accession Number:

BC012998

GeneID (NCBI):

10478

UNIPROT ID:

O43808

Full Name:

solute carrier family 25
(mitochondrial carrier; peroxisomal
membrane protein, 34kDa), member
17

Calculated MW:

307 aa, 35 kDa

Observed MW:

35 kDa

Purification Method:

Protein A purification

CloneNo.:

2D1D2

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:500-1:2000

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, Mouse

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : MCF-7 cells, U2OS cells, A549 cells, HeLa cells,
HepG2 cells, K-562 cells, Jurkat cells

IHC : human liver cancer tissue, human liver tissue,
mouse liver tissue

Background Information

SLC25A17 (solute carrier family 25 member 17), is a kind of peroxisomal transporter for multiple cofactors such as CoA, FAD, FMN and AMP. It may catalyze the transport of free CoA, FAD and NAD⁺ from the cytosol into the peroxisomal matrix by a counter-exchange mechanism. SLC25A17 is the only member of the mitochondrial carrier family localized to peroxisomal. (PMID: 22185573, 32266253)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67635-1-Ig (SLC25A17 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67635-1-Ig (SLC25A17 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).