## For Research Use Only

## **GSTT1** Monoclonal antibody

Size:

2200 µg/ml

Catalog Number: 67645-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number: 67645-1-lg BC007065

GeneID (NCBI): 2952

Source: UNIPROT ID:
Mouse P30711
Isotype: Full Name:

IgG2b glutathione S-transferase theta 1

Immunogen Catalog Number: Calculated MW:
AG8573 240 aa, 27 kDa

Observed MW: 28 kDa, 30 kDa

**Applications** 

Tested Applications: WB, IF-P, IHC, ELISA

Species Specificity: Human, pig, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed

with citrate buffer pH 6.0

Purification Method:

Protein A purification

CloneNo.: 1A4C11

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:250-1:1000 IF-P 1:200-1:800

Positive Controls:

WB: pig lung tissue, pig liver tissue, K-562 cells, rat

liver tissue, mouse brain tissue

IHC: mouse liver tissue,

IF-P: mouse liver tissue,

**Background Information** 

GSTT1 belongs to the GST superfamily and Theta family. It is a conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. GSTT1 acts on 1,2-epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. It displays glutathione peroxidase activity with cumene hydroperoxide. GSTM1 and GSTT1 has been suggested as a risk factor in various cancers. GSTT1 is important in the detoxification of unidentified xenobiotics in the large intestine.

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 67645-1-Ig (GSTT1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 67645-1-Ig (GSTT1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 67645-1-Ig (GSTT1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using GSTT1 antibody (67645-1-Ig, Clone: 1A4C11) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using GSTT1 antibody (67645-1-Ig, Clone: 1A4C11) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).