

For Research Use Only

NCAPH Monoclonal antibody

Catalog Number: 67655-1-Ig **Featured Product**



Basic Information

Catalog Number: 67655-1-Ig	GenBank Accession Number: BC024211	Purification Method: Protein A purification
Size: 700 µg/ml	GeneID (NCBI): 23397	CloneNo.: 3D2F11
Source: Mouse	UNIPROT ID: Q15003	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF 1:200-1:800
Isotype: IgG1	Full Name: non-SMC condensin I complex, subunit H	
Immunogen Catalog Number: AG27748	Calculated MW: 741 aa, 83 kDa	
	Observed MW: 83-100 kDa	

Applications

Tested Applications:

IF-P, IHC, WB, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, NIH/3T3 cells, Jurkat cells, K-562 cells, HSC-T6 cells, 4T1 cells

IHC: human liver cancer tissue, human colon cancer tissue

IF: human liver cancer tissue,

Background Information

Non-SMC condensin I complex subunit H (NCAPH) is one of the three non-SMC subunits in condensin I, which belongs to a recently defined superfamily of proteins termed kleisins. Another two non-SMC subunits, CAP-D2 and CAP-G, share a highly degenerate repeating motif known as HEAT repeat. Some studies show that each subunit is essential for viability and plays an important role in mitotic chromosome architecture and segregation. In recent years, researchers found that the high expression of NCAPH was associated with poor prognosis in patients with non-small cell lung cancer and prostate cancer. Downregulation of NCAPH inhibited the proliferation, migration, and invasion of several cancer cells significantly. Moreover, NCAPH was involved in the regulation of mature chromosome condensation and DNA damage. These data suggest that NCAPH may be a key carcinogen involved in the development and progression of human malignant tumors. (PMID: 28300828, PMID: 33311486)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67655-1-Ig (NCAPH antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67655-1-Ig (NCAPH antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using NCAPH antibody (67655-1-Ig, Clone: 3D2F11) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



WB result of NCAPH antibody (67655-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NCAPH transfected HEK-293 cells.