

For Research Use Only

SARS-CoV-2 Nucleocapsid Phosphoprotein Monoclonal antibody

Catalog Number: 67666-1-Ig **3 Publications**



Basic Information

Catalog Number:

67666-1-Ig

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG30676

GenBank Accession Number:

NC_045512

GeneID (NCBI):

43740575

Full Name:

COVID-19 N Protein

Purification Method:

Protein A purification

CloneNo.:

1B3C3

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

Virus

Cited Species:

mouse

Positive Controls:

WB : Ag30676,

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. It can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387). 67666-1-Ig can be used as capture antibody. 67666-2-Ig can be used as detection antibody.

Notable Publications

Author	Pubmed ID	Journal	Application
Marina Pribanić Matešić	35216036	Viruses	
I Novodchuk	35512584	Biosens Bioelectron	
Zhaohuan Wang	39287388	J Virol	

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 67666-1-Ig and 67666-2-Ig respectively. Signal was developed with TMB and stopped by H₂SO₄. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 67666-1-Ig at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRP-conjugated clone 67666-2-Ig was used at 1 μ g/mL for detection. Signal was developed with TMB and stopped by H₂SO₄. Signal strength was measured by absorbance at 450 nm.



E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 67666-1-Ig and 67666-2-Ig at various work concentration.