For Research Use Only

SARS-CoV-2 Nucleocapsid Phosphoprotein Monoclonal antibody



Catalog Number: 67666-2-lg

Basic Information

Catalog Number: 67666-2-Ig

Size: 1000 µg/ml Source:

Mouse Isotype: IgG2b

Immunogen Catalog Number:

AG30676

GenBank Accession Number:

NC_045512 GeneID (NCBI): 43740575 Full Name:

COVID-19 N Protein

Purification Method: Protein A purification

CloneNo.: 6D10E2

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications: WB,ELISA

Species Specificity:

Virus

Positive Controls:

WB: Ag30676,

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. it can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387).67666-1-lg can be used as capture antibody. 67666-2-lg can be used as detection antibody.


Storage

Storage: Store at -20°C. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 67666-1-lg and 67666-2-lg respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 67666-1-Ig at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRP-conjugated clone 67666-2-Ig was used at 1 μ g/mL for detection. Signal was developed with TMB and stopped by H2SO4 . Signal strength was measured by absorbance at 450 nm.



E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 67666-1-Ig and 67666-2-Ig at various work concentration.