For Research Use Only

LRPPRC Monoclonal antibody

Catalog Number:67679-1-lg

Featured Product

2 Publications



Basic Information

Catalog Number: GenBank Accession Number: 67679-1-lg BC050311 GeneID (NCBI): Concentration: 1000 ug/ml 10128 **UNIPROT ID:** Source: Mouse P42704 Full Name: Isotype:

lgG2b leucine-rich PPR-motif containing

Calculated MW: Immunogen Catalog Number: AG30261 1394 aa, 158 kDa Observed MW:

130 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IF

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

CloneNo.:

Purification Method:

Protein A purification

3H7E1

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:2000-1:8000 IF/ICC 1:200-1:800

Positive Controls:

WB: U2OS cells, U-251 cells, HeLa cells, HepG2 cells IHC: human colon cancer tissue, human endometrial cancer tissue, human liver cancer tissue

IF/ICC: HeLa cells, HEK-293 cells, HepG2 cells, A431

Background Information

LRPPRC (also called LRP130 or GP130) is a 130-kDa RNA-binding protein of the pentatricopeptide repeat family. LRPPRC localizes primarily to the mitochondria where it binds to poly(A) mRNA. It plays a role in translation or stability of mitochondrially encoded cytochrome c oxidase (COX) subunits. LRPPRC has also been shown to regulate $nuclear\ gene\ transcription\ and\ to\ bind\ specific\ RNA\ molecules\ in\ both\ the\ nucleus\ and\ the\ cytoplasm.\ Mutations\ in\ decreases and\ the\ cytoplasm.$ LRPPRC gene are associated with the French-Canadian type of Leigh syndrome.

Notable Publications

Author	Pubmed ID	Journal	Application
Dongqing Gu	38881895	Front Immunol	IF
Kejuan Jia	37139237	Front Genet	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67679-1-1g (LRPPRC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67679-1-Ig (LRPPRC antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67679-1-Ig (LRPPRC antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using LRPPRC antibody (67679-1-1g, Clone: 3H7E1) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using LRPPRC antibody (67679-1-lg, Clone: 3H7E1) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



WB result of LRPPRC antibody (67679-1-Ig; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-LRPPRC transfected HeLa cells.