For Research Use Only

DLD Monoclonal antibody

Catalog Number:67702-1-lg Featured Product

1 Publications

BC018696

1738

P09622

GeneID (NCBI):

UNIPROT ID:

Full Name:



Basic Information

Catalog Number: 67702-1-lg Concentration:

1000 ug/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG9680

dihydrolipoamide dehydrogenase Calculated MW: 509 aa, 54 kDa

GenBank Accession Number:

Observed MW: 54 kDa

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

IHC

Species Specificity: human, mouse, rat **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HuH-7 cells, NIH/3T3 cells, HSC-T6 cells, HeLa cells, LO2 cells, HepG2 cells, Jurkat cells

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:250-1:1000

CloneNo.:

2E7G5

IHC: human liver cancer tissue,

Background Information

DLD(Dihydrolipoyl dehydrogenase, mitochondrial) is also named as GCSL, LAD, PHE3 and belongs to the class-I pyridine nucleotide-disulfide oxidoreductase family. It catalyzes the oxidation of dihydrolipoamide, hE3 uses two molecules: non-covalently bound FAD and a transiently bound substrate, NAD+. DLD is involved in the hyperactivation of spermatazoa during capacitation and in the spermatazoal acrosome reaction.

Notable Publications

Author	Pubmed ID	Journal	Application
Kai Yao	37600770	Front Immunol	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67702-1-1g (DLD antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of DLD antibody (67702-1-lg; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DLD transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67702-1-Ig (DLD antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67702-1-1g (DLD antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).