For Research Use Only

TNC/Tenascin-C Monoclonal antibody, PBS Only

Size:



Purification Method:

Protein G purification

CloneNo.:

1F7G8

Catalog Number: 67710-1-PBS

Basic Information

Catalog Number: 67710-1-PBS

BC151843

GeneID (NCBI):

GenBank Accession Number:

1 mg/ml3371Source:UNIPROT ID:MouseP24821Isotype:Full Name:IgG1tenascin C

Immunogen Catalog Number: Calculated MW:

AG21412 2201 aa, 241 kDa
Observed MW:

260 kDa

Applications

Tested Applications:

WB,Indirect ELISA,IHC Species Specificity:

Human, Mouse, Rat

Background Information

Tenascin-C (TNC) is a large hexameric extracellular matrix glycoprotein of the tenascin family (PMID: 21818551). It is a multimodular protein containing multiple epidermal growth factor (EGF)-like repeats and fibronectin type III (FN III) domains (PMID: 1719530). Tenascin-C is highly expressed during embryonic development, particularly in the developing central nervous system, around motile cells and at epithelial-mesenchymal interaction sites (PMID: 25738825). In adult tissues, the expression and the distribution of TNC are typically limited under normal physiological conditions. It is upregulated during injury, inflammation, regeneration, and cancer (PMID: 7694605; 25120494). Tenascin-C is a diverse protein that can produce different functions including regulating cell adhesion, migration and proliferation.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67710-1-lg (TNC/Tenascin-C antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67710-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67710-1-1g (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67710-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67710-1-1g (TNC/Tenascin-C antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67710-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat brain tissue using TNC/Tenascin-C antibody (67710-1-1g, Clone: 1F7G8) at dilution of 1:200 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67710-1-PBS in a different storage buffer formulation.