For Research Use Only

CEP290 Monoclonal antibody

Catalog Number:67716-1-Ig




Basic Information	Catalog Number: 67716-1-lg	GenBank Accession Number: BC008641	Purification Method: Protein A purification	
	Size: 2100 µg/ml	GenelD (NCBI): 80184	CloneNo.: 2C10G2	
	Source: Mouse	UNIPROT ID: O15078	Recommended Dilutions: WB 1:1000-1:4000	
	lsotype: IgG2b	Full Name: centrosomal protein 290kDa		
	Immunogen Catalog Number: AG17457	Calculated MW: 2479 aa, 290 kDa		
		Observed MW: 290 kDa	IW:	
Applications	Tested Applications:	Positive C	Controls:	
	WB,ELISA Species Specificity: Human, rat, mouse		WB : HEK-293 cells, HSC-T6 cells, HeLa cells, NCCIT cells, K-562 cells	
		Studies in the Chlamydomonas model system has shown that the ciliary protein CEP290 is a critical component of these Y-link junctions. Additionally, numerous studies have demonstrated that CEP290's function is critical for IFT- in CEP290 knockdown experiments, many proteins that would normally localize to the cilium fail to do so, and cilium formation is disrupted or absent. Mutations in CEP290 are accountable for cases of nephronophthisis, Leber congenital amaurosis and Joubert syndrome. In IF analtsis of HeLa cells, the white arrows show centrosome and cilium staining. Two isoforms were produced by alternative splicing with predicted MW of 290 and 180 kDa according to UniProt.		
Background Information	these Y-link junctions. Additional in CEP290 knockdown experimen cilium formation is disrupted or a congenital amaurosis and Jouben cilium staining. Two isoforms we	ly, numerous studies have demonstra its, many proteins that would normally bsent. Mutations in CEP290 are accoun t syndrome. In IF analtsis of HeLa cells	ted that CEP290's function is critical for IFT / localize to the cilium fail to do so, and htable for cases of nephronophthisis, Leber s, the white arrows show centrosome and	
Background Informatio	these Y-link junctions. Additional in CEP290 knockdown experimen cilium formation is disrupted or a congenital amaurosis and Jouben cilium staining. Two isoforms we	ly, numerous studies have demonstra its, many proteins that would normally bsent. Mutations in CEP290 are accou t syndrome. In IF analtsis of HeLa cells re produced by alternative splicing wi	ted that CEP290's function is critical for IFT / localize to the cilium fail to do so, and htable for cases of nephronophthisis, Leber s, the white arrows show centrosome and	

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67716-1-1g (CEP290 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.