For Research Use Only

GATA2 Monoclonal antibody

Catalog Number:67722-1-Ig




Basic Information	Catalog Number: 67722-1-lg	GenBank Accession Number: BC018988	Purification Method: Protein A purification			
	Size: 2000 μg/ml	GenelD (NCBI): 2624	CloneNo.: 2B5A1			
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG1557	ENSEMBL Gene ID: ENSG00000179348 UNIPROT ID: P23769 Full Name: GATA binding protein 2	Recommended Dilutions: WB 1:1000-1:8000 IF/ICC 1:200-1:800			
				Observed MW: 50 kDa		
				Applications	Tested Applications:	Positive
		WB, IF/ICC, FC (Intra), ELISA Species Specificity: human, mouse, rat				HeLa cells, HUVEC cells, LNCaP cells, PC-12 cells 2 cells, HEK-293 cells, K-562 cells, NIH/3T3 cells 3 cells
IF/ICC : S	5H-SY5Y cells,					
Background Information	GATA2 is a member of GATA family of transcription factors that contains zinc fingers in DNA binding domain. GATA2 acts as a transcriptinal activator that regulates endothelin-1 expression in endothelial cells. Together with GATA3, GATA2 regulates adipocyte differentiation through molecular control of the preadpocyte-adipocyte transition. It also affect the activity of Rho inhibitor p190RhoGAP that controls capillary network formation in vitro and retinal angiogenesis in vivo.					
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20 ⁶	50% glycerol pH 7.3.				


 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 67722-1-Ig (GATA2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. 1X10^6 K-562 cells were intracellularly stained with 0.5 ug Anti-Human GATA2 (67722-1-Ig, Clone:2B5A1) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Mouse IgG1 (KFA022), or 0.5 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using GATA2 antibody (67722-1-Ig, Clone: 2B5A1) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).