For Research Use Only

ILK Monoclonal antibody

Catalog Number: 67724-1-lg 1 Publications



Basic Information

Catalog Number: 67724-1-lg

Size: 1000 ug/ml BC001554

GeneID (NCBI): 3611

GenBank Accession Number:

Source: UNIPROT ID:
Mouse Q13418
Isotype: Full Name:

IgG1 integrin-linked kinase

Immunogen Catalog Number:Calculated MW:AG21289451 aa, 59 kDa

Observed MW: 51 kDa Purification Method:

Protein G purification

CloneNo.: 1E10E4

Recommended Dilutions:

WB 1:2000-1:10000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, ELISA Cited Applications:

WB. IP

Species Specificity: human, mouse, rat, pig

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: U2OS cells, human placenta tissue, human peripheral blood platelets, rat colon tissue, mouse colon tissue, pig colon tissue, HepG2 cells, HeLa cells, HEK-293 cells. K-562 cells

IHC: human skin cancer tissue,

Background Information

ILK(Integrin-linked protein kinase) is also named as p59, PAT-4, DKFZp686F1765, ILK-1, ILK-2, ILK1, ILK2, p59ILK and belongs to the TKL Ser/Thr protein kinase family. This protein localizes to multiple sites, including the cytoplasm, and imports into the nucleus through sequences in its N-terminus, via active transport mechanisms that involve nuclear pore complexes(PMID: 18635968). ILK performs crucial roles in the control of intestinal cell and crypt-villus axis homeostasis-especially with regard to basement membrane fibronectin deposition-as well as cell proliferation, spreading, and migration(PMID:19885839).

Notable Publications

Author Pub	bmed ID J	ournal	Application
Yesheng Fu 385	582895 N	Nat Commun	WB,IP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67724-1-1g (ILK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffinembedded skin cancer slide using 67724-1-1g (ILK antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).