

For Research Use Only

IgG Kappa chain Monoclonal antibody



Catalog Number: 67761-1-Ig

Basic Information

Catalog Number:

67761-1-Ig

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Calculated MW:

155 kDa

Observed MW:

~25 kDa

Purification Method:

Protein A purification

CloneNo.:

4D11F7

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:2500-1:10000

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human plasma,

IHC : human tonsillitis tissue,

Background Information

This antibody detects the Kappa chain of human IgG,

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 67761-1-Ig (Human IgG Kappa chain antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 67761-1-Ig (Human IgG Kappa chain antibody) at dilution of 1:5000 (under 4x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 67761-1-Ig (Human IgG Kappa chain antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



16 nL human plasma were subjected to SDS PAGE followed by western blot with 67761-1-Ig (Human IgG Kappa chain antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.