

For Research Use Only

# CNOT4 Monoclonal antibody

Catalog Number: 67798-1-Ig **1 Publications**



## Basic Information

Catalog Number: 67798-1-Ig	GenBank Accession Number: BC035590	Purification Method: Protein A purification
Concentration: 1000 µg/ml	GeneID (NCBI): 4850	CloneNo.: 4H5H3
Source: Mouse	UNIPROT ID: O95628	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000 IF-P 1:200-1:800 IF/ICC 1:400-1:1600
Isotype: IgG2b	Full Name: CCR4-NOT transcription complex, subunit 4	
Immunogen Catalog Number: AG29964	Calculated MW: 575 aa, 64 kDa Observed MW: 64-85 kDa	

## Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls:
Species Specificity: human, mouse, rat	WB : HeLa cells, U2OS cells, MCF-7 cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells.
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	IHC : human colon cancer tissue, IF-P : human colon cancer tissue, IF/ICC : HepG2 cells,

## Background Information

CNOT4, also known as CCR4-NOT transcription complex subunit 4, has E3 ubiquitin ligase activity, and promotes ubiquitination and degradation of target proteins. CNOT4 is involved in activation of the JAK/STAT pathway. CNOT4 exist many isoforms and the range of its molecular weight from 48 to 84 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Jing Yang	39747175	Nat Commun	


## Storage

Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67798-1-Ig (CNOT4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.




Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using CNOT4 antibody (67798-1-Ig, Clone: 4H5H3 ) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).




WB result of CNOT4 antibody (67798-1-Ig; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CNOT4 transfected MCF-7 cells.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 67798-1-Ig (CNOT4 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using CNOT4 antibody (67798-1-Ig, Clone: 4H5H3 ) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CNOT4 antibody (67798-1-Ig, Clone: 4H5H3 ) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).