## For Research Use Only

## DAPK1 Monoclonal antibody

Catalog Number:67815-1-lg 3 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number: 67815-1-lg BC113660

Size: GeneID (NCBI): 1000  $\mu$  g/ml 1612

Source: UNIPROT ID: Mouse P53355
Isotype: Full Name:

lgG1 death-associated protein kinase 1

Immunogen Catalog Number: Calculated MW:

AG29838 1430 aa, 160 kDa Observed MW:

160 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, IF-P, FC (Intra), ELISA

**Cited Applications:** 

WB, IF

Species Specificity: human, mouse, rat Cited Species: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, A549 cells, HeLa cells, HepG2 cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, 4T1 cells

**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:1000-1:4000 IHC 1:200-1:800

IF-P 1:200-1:800

IF/ICC 1:400-1:1600

CloneNo.:

1E2F9

IHC: human breast cancer tissue, human placenta tissue, human stomach cancer tissue, mouse skin tissue, mouse small intestine tissue, rat small intestine tissue

IF-P: human breast cancer tissue,
IF/ICC: HCT 116 cells, HT-1376 cells

**Background Information** 

DAPK1(Death-associated protein kinase 1) is a stress-activated tumor suppressor protein that plays a role in both proapoptotic or antiapoptotic signal transduction pathways. Loss of DAPK1 expression is associated with a selective advantage fortumor cells to resist apoptotic stimuli, allowing them to separate from the original tumor; from this point of view, DAPK1 could be considered a tumor metastases inhibitor gene(PMID:17319784).

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Chao Geng	39479447	Theranostics	WB
Yunying Yang	39174646	Cell Death Differ	WB
Xiang-Xin Chen	37480108	Cell Commun Signal	IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 67815-1-Ig (DAPK1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67815-1-lg (DAPK1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using DAPK1 antibody (67815-1-lg, Clone: 1£2F9) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HT-1376 cells using DAPK1 antibody (67815-1-lg, Clone: 1E2F9) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



Immunofluorescent analysis of (-20°C Ethanol) fixed HCT 116 cells using DAPK1 antibody (67815-1-Ig, Clone: 1E2F9) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



1X10^6 HCT 116 cells were intracellularly stained with 0.4 ug Anti-Human DAPK1 (67815-1-lg, Clone:1E2F9) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).