For Research Use Only

DAPK1 Monoclonal antibody, PBS Only



Purification Method:

Protein G purification

CloneNo.:

1E2F9

Catalog Number: 67815-1-PBS

Basic Information

Catalog Number: GenBank Accession Number: BC113660

67815-1-PBS GeneID (NCBI): Size:

1mg/ml 1612 Source: **UNIPROT ID:** Mouse P53355 Isotype: Full Name:

lgG1 death-associated protein kinase 1

Calculated MW: Immunogen Catalog Number: AG29838 1430 aa, 160 kDa

> Observed MW: 160 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, Indirect ELISA

Species Specificity: human, mouse, rat

Background Information

DAPK1(Death-associated protein kinase 1) is a stress-activated tumor suppressor protein that plays a role in both proapoptotic or antiapoptotic signal transduction pathways. Loss of DAPK1 expression is associated with a selective advantage fortumor cells to resist apoptotic stimuli, allowing them to separate from the original tumor; from this point of view, DAPK1 could be considered a tumor metastases inhibitor gene(PMID:17319784).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67815-1-lg (DAPK1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 67815-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67815-1-lg (DAPK1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67815-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using DAPK1 antibody (67815-1-1g, Clone: 1E2F9) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+I). This data was developed using the same antibody clone with 67815-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HT-1376 cells using DAPK1 antibody (67815-1-lg, Clone: 1£2F9) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004). This data was developed using the same antibody clone with 67815-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HCT 116 cells using DAPK1 antibody (67815-1-lg, Clone: 1E2F9) at dilution of 1:800 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67815-1-PBS in a different storage buffer formulation.



1X10^6 HCT 116 cells were intracellularly stained with 0.4 ug Anti-Human DAPK1 (67815-1-Ig, Clone:1E2F9) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67815-1-PBS in a

