For Research Use Only

ID1 Monoclonal antibody

Catalog Number:67827-1-lg 4 Publications



Basic Information

Catalog Number: 67827-1-lg

Size: 1000 µg/ml Source: Mouse

Immunogen Catalog Number:

AG13359

Isotype:

lgG1

GenBank Accession Number:

3397

Calculated MW: 16 kDa Observed MW:

18 kDa

Applications

Tested Applications:

IHC, WB, ELISA **Cited Applications:**

WB. IHC

Species Specificity: Human, mouse, rat **Cited Species:** human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval be performed with aircraft buffer pH 6.0 with citrate buffer pH 6.0

BC000613 GeneID (NCBI):

UNIPROT ID: P41134

inhibitor of DNA binding 1, dominant negative helix-loop-helix protein

Positive Controls:

WB: A549 cells, HeLa cells, PC-3 cells, HepG2 cells,

Purification Method:

CloneNo.:

IHC 1:150-1:600

1H3G11

Protein G purification

Recommended Dilutions: WB 1:1000-1:6000

Jurkat cells, HSC-T6 cells, 4T1 cells IHC: human lung cancer tissue,

Background Information

ID (inhibitor of DNA binding) proteins contain a helix-loop-helix (HLH) motif and regulate tissue-specific transcription within several cell lineages. They do not bind DNA directly, but inhibit lineage commitment by binding basic helix-loop-helix (bHLH) transcription factors through their HLH motif. ID1 proteins lack a basic DNAbinding domain but are able to form heterodimers with other HLH proteins, thereby inhibiting DNA binding. ID proteins contribute to cell growth, senescence, differentiation, and angiogenesis. Inhibitor of differentiation or DNA $binding - 1 \ (ID1) \ interacts \ with the \ transactivation \ domain \ of \ p65 \ (p65 TAD) \ both \ in \ vitro \ and \ in \ vivo. \ The \ molecular$ weight of ID1 is about 18 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Lijun Li	34658848	Front Pharmacol	WB
Lin Lu	35487890	Oncogenesis	WB
Zimei Zeng	37416767	Int J Biol Sci	WB,IHC

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67827-1-1g (ID1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 67827-1-1g (ID1 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).