For Research Use Only

TARS Monoclonal antibody

Catalog Number:67828-1-lg 1 Publications




GenBank Accession Number:Purification Method:BC000517Protein A purification
GeneID (NCBI): CloneNo.: 6897 1A11F10
68971A11F10UNIPROT ID:Recommended Dilutions:P26639WB 1:5000-1:50000
Full Name: IF 1:400-1:1600 threonyl-tRNA synthetase
er: Calculated MW: 83 kDa
Observed MW: 83-87 kDa
Positive Controls:
WB : HeLa cells, HSC-T6 cells, PC-12 cells, 4T1 cells HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells
IF : HeLa cells,
stase), also known as threonine-tRNA ligase or ThrRS, belongs to the aminoacyl-tRNA yl-tRNA synthetase (aaRS) catalyzes the specific esterification of an amino acid witl of its cognate tRNA. Specificity of aminoacylation catalyzed by aaRSs ensures the enetic code in the first step of protein synthesis (PMID: 26324710). TARS has some mass of 83, 87 kDa and other MW.
Pubmed ID Journal Application
35659036 J Hematol Oncol WB

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67828-1-Ig (TARS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67828-1-Ig (TARS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. 4T1 cells were subjected to SDS PAGE followed by western blot with 67828-1-1g (TARS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

150 kDa-


100 kDa

70 kDa

50 kDa-

40 kDa-

30 kDa→



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using TARS antibody (67828-1-Ig, Clone: 1A11F10) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).