

For Research Use Only

BAD Monoclonal antibody

Catalog Number: 67830-1-Ig **3 Publications**



Basic Information

Catalog Number:

67830-1-Ig

Size:

1000 μ g/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

BC001901

GeneID (NCBI):

572

UNIPROT ID:

Q92934

Full Name:

BCL2-associated agonist of cell death

Calculated MW:

18 kDa

Observed MW:

18-25 kDa

Purification Method:

Protein A purification

CloneNo.:

4B12G2

Recommended Dilutions:

WB 1:1000-1:6000

IF 1:200-1:800

Applications

Tested Applications:

IF/ICC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human, Mouse, Rat

Cited Species:

human, mouse, rat

Positive Controls:

WB : LNCaP cells, RAW 264.7 cells, HeLa cells, HEK-293 cells, Jurkat cells, Raji cells, THP-1 cells, HSC-T6 cells

IF : HeLa cells,

Background Information

BAD, also named as BBC6 and BCL2L8, promotes cell death. It is successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W, thereby affecting the level of heterodimerization of these proteins with BAX. BAD can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. It appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

Notable Publications

Author	Pubmed ID	Journal	Application
Yaling Zhang	36062194	Oxid Med Cell Longev	WB
Xueying Chen	33817830	FASEB J	WB
Zhong Xian	37062136	Phytomedicine	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67830-1-Ig (BAD antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using BAD antibody (67830-1-Ig, Clone: 4B12G2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).