

For Research Use Only

OXCT1 Monoclonal antibody, PBS Only



Catalog Number: 67836-1-PBS

Featured Product

Basic Information

Catalog Number:

67836-1-PBS

Size:

1mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG24792

GenBank Accession Number:

BC009001

GeneID (NCBI):

5019

UNIPROT ID:

P55809

Full Name:

3-oxoacid CoA transferase 1

Calculated MW:

520 aa, 56 kDa

Observed MW:

52-56 kDa

Purification Method:

Protein G purification

CloneNo.:

1G1B9

Applications

Tested Applications:

Indirect ELISA, IF/ICC, IHC, WB

Species Specificity:

rabbit, pig, rat, mouse, human

Background Information

OXCT1 (also known as SCOT), encoded by nuclear gene, is a mitochondrial CoA transferase required for ketone body degradation. It catalyzes the transfer of CoA from succinyl-CoA to acetoacetate, generating acetoacetyl-CoA. OXCT1 is expressed in brain, heart, and skeletal muscle, but not in liver. This antibody specifically recognizes endogenous OXCT1. (21209089)

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




rat heart tissue were subjected to SDS PAGE followed by western blot with 67836-1-Ig (SCOT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.




WB result of OXCT1 antibody (67836-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-OXCT1 transfected HeLa cells. This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using OXCT1 antibody (67836-1-Ig, Clone: 1G1B9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). CL594-Phalloidin (red). This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 67836-1-Ig (OXCT1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OXCT1 antibody (67836-1-Ig, Clone: 1G1B9) at dilution of 1:500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using OXCT1 antibody (67836-1-Ig, Clone: 1G1B9) at dilution of 1:500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 67836-1-PBS in a different storage buffer formulation.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human OXCT1 (67836-1-Ig, Clone:1G1B9) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67836-1-PBS in a

