For Research Use Only

PSMB1 Monoclonal antibody, PBS Only

Catalog Number: 67847-1-PBS



Purification Method:

Protein G purification

CloneNo.:

1C7C5

Basic Information

Catalog Number: 67847-1-PBS

Concentration: 1mg/ml

BC020807 GeneID (NCBI): 5689

GenBank Accession Number:

Source: UNIPROT ID:
Mouse P20618
Isotype: Full Name:

lgG1 proteasome (prosome, macropain)
subunit, beta type, 1

Immunogen Catalog Number: subunit, beta type,
AG30791 Calculated MW:
241 aa, 26 kDa

Observed MW: 25 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), Indirect ELISA

Species Specificity:

human

Background Information

PSMB1(Proteasome subunit beta type-1) is also named as PSC5 and belongs to the peptidase T1B family. The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The gene encodes a 241 amino acid protein with a 28 amino acid propeptide and two glycosylation sites.

Storage


Storage:

Store at -80°C


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67847-1-lg (PSMB1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67847-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67847-1-lg (PSMB1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67847-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using P5MB1 antibody (67847-1-1g, Clone: 1C7C5) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67847-1-PBS in a different storage buffer formulation.



1x10^6 Jurkat cells were intracellularly stained with 0.25 ug PSMB1 Monoclonal antibody (67847-1-1g, Clone:1C7C5) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (5A00013-1)(red), or 0.25 ug Mouse IgG1 isotype control Mouse McAb (66360-1-1g, Clone: 1F8D3) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67847-1-PBS in a different storage buffer