For Research Use Only

AFP Monoclonal antibody, PBS Only

Catalog Number:67852-1-PBS




Basic Information	Catalog Number: 67852-1-PBS	GenBank Accession Number: BC027881	Purification Method: Protein A purification
	Size:	GeneID (NCBI):	CloneNo.: 4B3E4
	1mg/ml Source: Mouse	174 UNIPROT ID: P02771	4DJE4
	lsotype: IgG1	Full Name: alpha-fetoprotein	
	Immunogen Catalog Number: AG6089	Calculated MW: 69 kDa	
		Observed MW: 68-72 kDa	
Applications	Tested Applications: WB, Indirect ELISA		
	Species Specificity: Human		
Background Information	AFP, also known as Alpha-fetoprotein, is a major plasma protein in the fetus and its concentration is very low in the adult (PubMed:24120489). AFP is considered a marker of hepatocyte precursors. AFP can be detected at abnormally high concentrations in hepatocellular carcinomas as well as in the plasma and ascitic fluid of adults with hepatoma High AFP concentrations have been correlated with tumor cell growth, indicating that AFP can serve as a tumor marker (PubMed: 18669658).		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	yat-80°C


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 67852-1-1g (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67852-1-PBS in a different storage buffer formulation.



HuH-7 cells were subjected to SDS PAGE followed by western blot with 67852-1-1g (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67852-1-PBS in a different storage buffer formulation.




human testis tissue were subjected to SDS PAGE followed by western blot with 67852-1-Ig (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67852-1-PBS in a different storage buffer formulation.




human placenta tissue were subjected to SDS PAGE followed by western blot with 67852-1-lg (AFP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed

using the same antibody clone with 67852-1-PBS in a different storage buffer formulation.



formulation.

Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67852-1-1g (AFP antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67852-1-PBS in a different storage buffer



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 67852-1-Ig (AFP antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67852-1-PBS in a different storage buffer formulation.