

Phospho-S6 Ribosomal protein (Ser235) Monoclonal antibody

 Catalog Number: 67898-1-Ig **3 Publications**

Basic Information

Catalog Number:

67898-1-Ig

Concentration:

500 ug/ml

Source:

Mouse

Isotype:

IgG2b

GenBank Accession Number:

BC000524

GeneID (NCBI):

6194

UNIPROT ID:

P62753

Full Name:

ribosomal protein S6

Calculated MW:

29 kDa

Observed MW:

32 kDa

Purification Method:

Protein A purification

CloneNo.:

2A4B6

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:1000-1:4000

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : HeLa cells, HEK-293 cells, Calyculin A treated HeLa cells, Calyculin A treated HEK-293 cells, IGF-1 treated MCF-7 cells, Calyculin A treated NIH/3T3 cells

IF/ICC : 100 nM Calyculin A (30 minutes) treated HeLa cells,

Background Information

Ribosomal protein S6 (RPS6) is one of the components of the 40S ribosomal subunit. RPS6 has been functionally regarded as the stimulator and/or inhibitor of certain types of mRNA translation, as well as the regulator of cellular metabolisms, cells size, survival and proliferation. RPS6 is phosphorylated at multiple sites, comprised between Ser235 and Ser247, by the p70 rpS6 kinase (S6K) 1, which is a major downstream effector of the mammalian target of rapamycin complex 1 (mTORC1). Phosphorylation of RPS6 at the dual site Ser235/236 occurs also independently of mTORC1, via the p90 ribosomal S6 kinases (RSK), which are activated by the extracellular signal-regulated kinases (ERK). Recent studies performed in pancreatic β -cells identified PKA as an additional RPS6 kinase, specifically involved in the phosphorylation of Ser235/236. (PMID: 26490682, PMID: 21814187, PMID: 31112404). 67898-1-Ig specifically recognizes the phosphorylation site of Ser235 or dual site Ser235/236.

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Tao	31908053	FASEB J	WB
Evangelia Lekka	36572670	Nat Commun	WB
Joao Paulo Cavalcanti de Albuquerque	40184437	Sci Immunol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 67898-1-Ig (Phospho-S6 Ribosomal protein (Ser235) antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta tubulin antibody as loading control.



Immunofluorescent analysis of (4% PFA) fixed untreated and 100 nM Calyculin A (30 minutes) treated HeLa cells using Phospho-S6 Ribosomal protein (Ser235) antibody (67898-1-Ig, Clone: 2A4B6) at dilution of 1:2000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



1×10^6 untreated HeLa cells (dash lines) and 100 nM Calyculin A (30 minutes) treated HeLa cells (full lines) were intracellularly stained with $0.2 \mu\text{g}$ Phospho-S6 Ribosomal protein (Ser235) Monoclonal antibody (67898-1-Ig, Clone: 2A4B6, red) and Multi-rAb CoraLite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM005). Mouse IgG2b isotype control Mouse McAb (66360-3-Ig, Clone: 11B8C4, blue) was parallel stained as control. Cells were fixed