For Research Use Only

Phospho-S6 Ribosomal protein (Ser235) Monoclonal antibody



Catalog Number:67898-1-lg 2 Publications

Basic Information

Catalog Number: 67898-1-lg Size: 500 μg/ml

Source: Mouse Isotype:

lgG2b

GenBank Accession Number: BC000524

GeneID (NCBI): 6194 **UNIPROT ID:**

P62753 Full Name:

32 kDa

ribosomal protein S6 Calculated MW: 29 kDa Observed MW:

Purification Method:

Protein A purification

CloneNo.: 2A4B6

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: Human, mouse, rat Cited Species: human

Positive Controls:

WB: HeLa cells, HEK-293 cells, Calyculin A treated HeLa cells, Calyculin A treated HEK-293 cells, IGF-1 treated MCF-7 cells, Calyculin A treated NIH/3T3 cells

Background Information

Ribosomal protein S6 (RPS6) is one of the components of the 40S ribosomal subunit. RPS6 has been functionally regarded as the stimulator and/or inhibitor of certain types of mRNA translation, as well as the regulator of cellular metabolisms, cells size, survival and proliferation. RPS6 is phosphorylated at multiple sites, comprised between Ser235 and Ser247, by the p70 rpS6 kinase (S6K) 1, which is a major downstream effector of the mammalian target of rapamycin complex 1 (mTORC1). Phosphorylation of RPS6 at the dual site Ser235/236 occurs also independently of mTORC1, via the p90 ribosomal S6 kinases (RSK), which are activated by the extracellular signal-regulated kinases (ERK). Recent studies performed in pancreatic β-cells identified PKA as an additional RPS6 kinase, specifically involved in the phosphorylation of Ser235/236. (PMID: 26490682, PMID: 21814187, PMID: 31112404). $67898 \hbox{-} 1-lg\ specifically\ recognizes\ the\ phosphorylation\ site\ of\ Ser 235\ or\ dual\ site\ Ser 235/236.$

Notable Publications

Author	Pubmed ID	Journal	Application
Yu Tao	31908053	FASEBJ	WB
Evangelia Lekka	36572670	Nat Commun	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 67898-1-Ig (Phospho-56 Ribosomal protein (Ser235) antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta tubulin antibody as loading control.