

For Research Use Only

HDAC11 Monoclonal antibody, PBS Only



Catalog Number: 67949-1-PBS

Basic Information

Catalog Number: 67949-1-PBS	GenBank Accession Number: BC009676	Purification Method: Protein G purification
Size: 1 mg/ml	GeneID (NCBI): 79885	CloneNo.: 1G12B5
Source: Mouse	UNIPROT ID: Q96DB2	
Isotype: IgG1	Full Name: histone deacetylase 11	
Immunogen Catalog Number: AG28335	Calculated MW: 347 aa, 39 kDa	
	Observed MW: 39 kDa	

Applications

Tested Applications:
WB, Indirect ELISA, IHC

Species Specificity:
Human

Background Information

Histone deacetylases (HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. At least 4 classes of HDAC were identified. HDAC11 belongs to Class IV HDACs. HDAC11 is localized to the nucleus and may be involved in regulating the expression of interleukin 10. Alternative splicing results in multiple transcript variants. HDAC11 has 2 isoforms with the molecular mass of 33 and 39 kDa.

Storage

Storage:
Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67949-1-Ig (HDAC11 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67949-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 67949-1-Ig (HDAC11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67949-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 67949-1-Ig (HDAC11 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67949-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 67949-1-Ig (HDAC11 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67949-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 67949-1-Ig (HDAC11 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67949-1-PBS in a different storage buffer formulation.