## For Research Use Only

## Caspase 7 Monoclonal antibody

Catalog Number:67956-1-lg 2 Publications





Basic Information	Catalog Number: 67956-1-lg	GenBank Accession Number: BC015799		Purification Method: Protein A purification	
	Size:	GeneID (NCBI):		CloneNo.:	
	1000 µg/ml	g/ml 840		3C9H4	
	Source: UNIPROT ID:			Recommended Dilutions:	
	Mouse	P55210		WB 1:5000-1:50000 IHC 1:500-1:2000	
	Isotype: IgG2a	Full Name:	caspase 7, apoptosis-related cysteine		
	Immunogen Catalog Number: AG27601	peptidase Calculated MW:			
					303 aa, 34 kDa
			Observed MW:		
		35 kDa			
	Applications	Tested Applications:	ted Applications: Positive Controls:		rols:
WB, IHC, ELISA		WB : Jurkat cells, PC-12 cells, RAW 264.7 cells			
Cited Applications: WB, IF		IHC : rat stomach tissue, h		ach tissue, human liver cancer tissue,	
Species Specificity:		mouse liver tissue, mouse stomach tiss		issue, mouse stomach tissue, rat liver	
Species Specificity: tissue   Human, mouse, rat tissue					
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0					
Background Information	Caspase 7(CASP7), like caspases 3 and 6, contains a short prodomain and, upon apoptotic induction, the 35 kDa proform is converted into a 32 kDa intermediate or preactive form which is further processed into two active subunits consisting of the p20 or large (18 kDa) subunit and the p10 or small (11 kDa) subunit and it is present in the brain, which is up-regulated and activated after traumatic injury(PMID:15953353). Caspase-7 is classified as a member of the subgroup of cysteine proteases most related to the Caenorhabditis elegans factor CED-3, which also includes caspase-3, -6, and -9(PMID:9426061). The protein is involved in the activation cascade of caspases responsible for apoptosis execution.				
Notable Publications	Author	Pubmed ID Journ	al	Application	
	Shunze Jia	39460804 Cell M	1ol Life Sci	WB	
	Qi Sun	39277993 Aquat	Toxicol	IF	


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data








Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67956-1-1g (Caspase 7 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67956-1-1g (Caspase 7 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 67956-1-Ig (Caspase 7 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse stomach tissue slide using 67956-1-1g (Caspase 7 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded rat liver tissue slide using 67956-1-lg (Caspase 7 antibody) at dilution of 1:1000 (under 10x (ens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).





Immunohistochemical analysis of paraffinembedded rat stomach tissue slide using 67956-1-Ig (Caspase 7 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffinembedded rat stomach tissue slide using 67956-1-Ig (Caspase 7 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).