For Research Use Only

PSMB5 Monoclonal antibody, PBS Only

Catalog Number:67959-1-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

2E9H1

Basic Information

Catalog Number: 67959-1-PBS Concentration:

1mg/ml
Source:
Mouse

IgG2a Immunogen Catalog Number:

AG5828

AG5828

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), Indirect ELISA

Species Specificity:

human

Isotype:

Background Information

PSMB5(Proteasome subunit beta type-5) is also named as LMPX, MB1, X and belongs to the peptidase T1B family. The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH.RNAi-mediated silencing of. It has 2 isoforms produced by alternative splicing.

GenBank Accession Number:

proteasome (prosome, macropain)

BC057840

5693

P28074

GeneID (NCBI):

UNIPROT ID:

Full Name:

subunit, beta type, 5

Calculated MW: 263 aa, 28 kDa Observed MW: 22 kDa

Storage

Applications


Storage:

Store at -80°C

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67959-1-1g (PSMB5 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67959-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67959-1-Ig (PSMB5 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67959-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PSMB5 antibody (67959-1-lg, Clone: 2E9H1) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+I). This data was developed using the same antibody clone with 67959-1-PBS in a different storage buffer formulation.



WB result of PSMB5 antibody (67959-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PSMB5 transfected HeLa cells. This data was developed using the same antibody clone with 67959-1-PBS in a different storage buffer formulation.



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug PSMB5 Monoclonal antibody (67959-1-1g, Clone:2E9H1) and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG2a isotype control Mouse McAb (66360-2-1g, Clone: 11A1B2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 67959-1-PBS in a different storage buffer