For Research Use Only

PRMT6 Monoclonal antibody

Catalog Number:67981-1-lg




Basic Information	Catalog Number: 67981-1-lg	GenBank Accession Number: BC002729	Purification Method: Protein A purification	
	Size: 1000 µg/ml	GenelD (NCBI): 55170	CloneNo.: 4B5A9	
	Source: Mouse	UNIPROT ID: Q96LA8	Recommended Dilutions: WB 1:5000-1:50000	
	lsotype: IgG2a	Full Name: protein arginine methyltransferase 6 ber: Calculated MW: 42 kDa		
	Immunogen Catalog Number: AG7934			
		Observed MW: 42 kDa		
Applications	Tested Applications:	Positive	Positive Controls:	
	WB, ELISA Species Specificity: Human, Mouse, Rat	WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells		
Background Information	Protein arginine methyltransferase 6 (PRMT6) is a type I PRMT which is involved in epigenetic regulation of gene expression, alternative splicing, development and differentiation, DNA repair, cell proliferation and senescence, DNA methylation, mitosis, inflammation, innate antiviral immunity, spermatogenesis, transactivation of nuclear receptors and cell signaling. The human PRMT6 gene, located on Chromosome 1, encodes for the 41.9 kDa PRMT6 enzyme. PRMT6 is predominantly localized to the nucleus, in stark contrast to PRMT3 and PRMT5 which are preponderantly cytosolic, while other PRMTs are found in both nucleus and cytosol. PRMT6 is expressed in a wide range of tissues with high expression in kidney and testes. PRMT6 generates asymmetric dimethylation modifications in histone 3 at arginine 2, arginine 17 and arginine 42 (H3R2me2a, H3R17me2a and H3R42me2a) and in histone H2A at arginine 26 (H2AR26me2a) and participates in the epigenetic regulation of gene expression.			
	in histone H2A at arginine 26 (H2A		-	

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67981-1-Ig (PRMT6 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.