For Research Use Only

ACOX1 Monoclonal antibody

Catalog Number:68017-1-lg Featured Product

3 Publications

BC010425



Basic Information

Catalog Number: 68017-1-lg Size:

1000 µg/ml Source: Mouse

Isotype: lgG1

Immunogen Catalog Number:

AG28037

UNIPROT ID:

Q15067 Full Name:

GeneID (NCBI):

acyl-Coenzyme A oxidase 1, palmitovl

GenBank Accession Number:

Calculated MW: 74 kDa Observed MW:

50 kDa

Purification Method:

Protein A purification

CloneNo.: 3E12E10

Recommended Dilutions: WB 1:2000-1:10000

Applications

Tested Applications: WB, ELISA

Cited Applications: WB, IF, IHC

Species Specificity: Human, Rat, Mouse, pig **Cited Species:** human, rat, mouse

Positive Controls:

WB: LNCaP cells, pig liver tissue, rat liver tissue, HepG2 cells, HeLa cells, HEK-293 cells, Jurkat cells,

HSC-T6 cells, NIH/3T3 cells

Background Information

ACOX1(Peroxisomal acyl-coenzyme A oxidase 1) is the first and rate-limiting enzyme in the peroxisomal fatty acid beta-oxidation pathway and catalyzes the desaturation of acyl-CoAs to 2-trans-enoyl-CoAs. It belongs to the acyl-CoA oxidase family and also named as ACOX, AOX, SCOX. It has 3 isoforms produced by alternative splicing with the molecular mass of 70-74 kDa and can be cleaved post-translationally into a 50-kDa and a 22-kDa subunit between Val468 and Ala469 (PMID: 10672038). Defects in ACOX1 are the cause of adrenoleukodystrophy pseudoneonatal (Pseudo-NALD).

Notable Publications

Author	Pubmed ID	Journal	Application
Xinhui Liu	36313325	Front Pharmacol	WB
Qiming Xiao	35567992	Phytomedicine	WB,IF
Yang-He Zhang	38367917	Pharmacol Res	WB,IHC,IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68017-1-lg (ACOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



rat liver tissue were subjected to SDS PAGE followed by western blot with 68017-1-1g (ACOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of ACOX1 antibody (68017-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACOX1 transfected HepG2 cells.



pig liver tissue were subjected to SDS PAGE followed by western blot with 68017-1-lg (ACOX1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.