

FXVD6 Monoclonal antibody

Catalog Number: 68058-1-Ig

Basic Information

Catalog Number: 68058-1-Ig	GenBank Accession Number: BC018652	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 53826	CloneNo.: 2H1B7
Source: Mouse	UNIPROT ID: Q9HQ3	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:400-1:1600
Isotype: IgG1	Full Name: FXVD domain containing ion transport regulator 6	
Immunogen Catalog Number: AG8538	Calculated MW: 95 aa, 11 kDa	
	Observed MW: 20 kDa	

Applications

Tested Applications:
IF/ICC, IHC, WB, ELISA

Species Specificity:
Human, Mouse, Rat, Pig, Rabbit, Chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: PC-12 cells, rat cerebellum tissue, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue, mouse cerebellum tissue

IHC: mouse cerebellum tissue, mouse brain tissue

IF/ICC: PC-12 cells,

Background Information

The FXVD family is a group of small single-span transmembrane proteins characterized by a signature sequence containing an FXVD motif, two conserved glycines and a serine residue. Members of the FXVD family, including FXVD1 (phospholemmann), FXVD2 (gamma subunit of Na,K-ATPase), FXVD3 (Mat8), FXVD4 (CHIF), FXVD5 (RIC), FXVD6 (phosphohippolin) and FXVD7, are tissue specific regulators of the Na,K-ATPase. FXVD6 is primarily expressed in the brain. It modulates the kinetic activity of Na,K-ATPase and has long-term physiological importance in maintaining cation homeostasis. It may play a role in endolymph composition and has a potential important role in neuronal excitability of the CNS during postnatal development and in the adult brain. On the SDS-PAGE FXVD6 migrates with an apparent molecular weight of approximately 20 kDa, which is larger than the calculated molecular weight of 10.5 kDa (PMID: 15193427; 17209044). The gene encodes FXVD6 is located on chromosome 11q23.3, and it might be a susceptibility gene of schizophrenia.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68058-1-Ig (FXYD6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 68058-1-Ig (FXYD6 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 68058-1-Ig (FXYD6 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed PC-12 cells using FXYD6 antibody (68058-1-Ig, Clone: 2H1B7) at dilution of 1:800 and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).