

# FXVD6 Monoclonal antibody

Catalog Number: 68058-1-Ig

## Basic Information

<b>Catalog Number:</b> 68058-1-Ig	<b>GenBank Accession Number:</b> BC018652	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 53826	<b>CloneNo.:</b> 2H1B7
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q9H0Q3	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IHC 1:500-1:2000 IF 1:400-1:1600
<b>Isotype:</b> IgG1	<b>Full Name:</b> FXVD domain containing ion transport regulator 6	
<b>Immunogen Catalog Number:</b> AG8538	<b>Calculated MW:</b> 95 aa, 11 kDa	
	<b>Observed MW:</b> 20 kDa	

## Applications

**Tested Applications:**  
IF/ICC, IHC, WB, ELISA

**Species Specificity:**  
Human, Mouse, Rat, Pig, Rabbit, Chicken

**Note-IHC:** suggested antigen retrieval with **TE buffer pH 9.0**; (\*) Alternatively, antigen retrieval may be performed with **citrate buffer pH 6.0**

**Positive Controls:**

**WB:** PC-12 cells, rat cerebellum tissue, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue, mouse cerebellum tissue

**IHC:** mouse cerebellum tissue, mouse brain tissue

**IF:** PC-12 cells,

## Background Information

The FXVD family is a group of small single-span transmembrane proteins characterized by a signature sequence containing an FXVD motif, two conserved glycines and a serine residue. Members of the FXVD family, including FXVD1 (phospholemmann), FXVD2 (gamma subunit of Na,K-ATPase), FXVD3 (Mat8), FXVD4 (CHIF), FXVD5 (RIC), FXVD6 (phosphohippolin) and FXVD7, are tissue specific regulators of the Na,K-ATPase. FXVD6 is primarily expressed in the brain. It modulates the kinetic activity of Na,K-ATPase and has long-term physiological importance in maintaining cation homeostasis. It may play a role in endolymph composition and has a potential important role in neuronal excitability of the CNS during postnatal development and in the adult brain. On the SDS-PAGE FXVD6 migrates with an apparent molecular weight of approximately 20 kDa, which is larger than the calculated molecular weight of 10.5 kDa (PMID: 15193427; 17209044). The gene encodes FXVD6 is located on chromosome 11q23.3, and it might be a susceptibility gene of schizophrenia.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68058-1-Ig (FXYD6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 68058-1-Ig (FXYD6 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 68058-1-Ig (FXYD6 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed PC-12 cells using FXYD6 antibody (68058-1-Ig, Clone: 2H1B7 ) at dilution of 1:800 and CoraLite@594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).