

For Research Use Only

INO80 Monoclonal antibody

Catalog Number: 68064-1-Ig



Basic Information

Catalog Number: 68064-1-Ig	GenBank Accession Number: BC146785	Purification Method: Protein G purification
Size: 1500 µg/ml	GeneID (NCBI): 54617	CloneNo.: 1F11C5
Source: Mouse	UNIPROT ID: Q9ULG1	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:250-1:1000
Isotype: IgG1	Full Name: INO80 homolog (S. cerevisiae)	
Immunogen Catalog Number: AG20265	Calculated MW: 1556 aa, 177 kDa	
	Observed MW: 177 kDa	

Applications

Tested Applications:

FC, IHC, WB, ELISA

Species Specificity:

Human, Rat, Mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : K-562 cells, HeLa cells, HEK-293 cells, Jurkat cells, NIH/3T3 cells, RAW 264.7 cells, PC-12 cells

IHC : human colon cancer tissue,

Background Information

DNA helicase INO80, also named as INO80 or hINO80, is a 1,556 amino acid protein, which contains one helicase C-terminal domain, one helicase ATP-binding domain, and one DBINO domain. INO80 co-localizes with PCMA at replication forks during S-phase in the nucleus and belongs to the SNF2 helicase family. INO80 is a component of the chromatin remodeling INO80 complex and is widely expressed in many tissues, such as brain, liver and pancreas. INO80 is involved in transcription regulation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68064-1-Ig (INO80 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 68064-1-Ig (INO80 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



1X10⁶ PC-12 cells were intracellularly stained with 0.8 ug Anti-Human INO80 (68064-1-Ig, Clone:1F11C5) and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.8 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 68064-1-Ig (INO80 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 68064-1-Ig (INO80 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).