## For Research Use Only

# BAG3 Monoclonal antibody

Catalog Number:68076-1-lg 2 Publications



**Basic Information** 

Catalog Number: 68076-1-lg

Size: 1000 µg/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG17608

Observed MW: 74 kDa

Positive Controls:

WB: NIH/3T3 cells, LNCaP cells, rabbit heart tissue,

**Purification Method:** 

CloneNo.:

4G6C2

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000 IF 1:500-1:2000

HeLa cells, pig heart, rat heart

IF: HepG2 cells,

**Applications Tested Applications:** 

IF/ICC, WB, ELISA Cited Applications: WB, IF, IHC Species Specificity:

Human, mouse, rabbit, pig, rat

**Cited Species:** human, rat

## **Background Information**

BAG3 (Bcl2-associated athanogene 3) belongs to the BAG protein family, the co-chaperone that binds to Hsc70/Hsp70 through the BAG domain and modulates their activity in polypeptide folding. BAG3 contains also a WW domain and a proline-rich (PXXP) repeat, that mediate binding to partners different from Hsp70. Through  $interacting\ with\ different\ molecular\ partner,\ BAG3\ influences\ several\ cell\ processes,\ such\ as\ apoptosis,\ autophagy$ and cell motility. BAG3 protein has been reported to sustain cell survival, resistance to therapy, and/or motility and metastatization in several tumor types, thus being identified as a potential target for anticancer therapies. In addition, defects in BAG3 are the cause of some myopathy.

GenBank Accession Number:

BCL2-associated athanogene 3

BC006418

9531

095817

61 kDa

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

Calculated MW:

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xin Lu	35253089	J Physiol Biochem	IHC
Jiao Lu	37990150	J Physiol Sci	WB,IF


Storage

Storage: Store at -20°C.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

#### **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 68076-1-1g (BAG3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using BAG3 antibody (68076-1-Ig, Clone: 4G6C2) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).