

For Research Use Only

TACSTD2/TROP2 Monoclonal antibody

Catalog Number: 68141-1-Ig



Basic Information

Catalog Number: 68141-1-Ig	GenBank Accession Number: BC009409	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 4070	CloneNo.: 5A6B1
Source: Mouse	UNIPROT ID: P09758	Recommended Dilutions: WB 1:1000-1:5000 IHC 1:500-1:2000 IF/ICC 1:400-1:1600
Isotype: IgG2b	Full Name: tumor-associated calcium signal transducer 2	
Immunogen Catalog Number: AG25775	Calculated MW: 36 kDa	
	Observed MW: 40-50 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls:
Species Specificity: human	WB : HaCaT cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human breast cancer tissue, IF/ICC : HaCaT cells,

Background Information

Trophoblast cell surface antigen 2 (TROP2), also named as TACSTD2, is a transmembrane glycoprotein. TROP2 plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells, cornea and multi-stratified epithelia. TROP2 is overexpressed in most carcinomas and is involved in cancer proliferation, migration, invasion, and metastasis. Congenital mutations of TROP2 cause a rare autosomal recessive disease which may lead to blindness-the gelatinous drop-like corneal disease (PMID: 35688908).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HaCaT cells were subjected to SDS PAGE followed by western blot with 68141-1-Ig (TACSTD2/TROP2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 68141-1-Ig (TACSTD2/TROP2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 68141-1-Ig (TACSTD2/TROP2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HaCaT cells using TACSTD2/TROP2 antibody (68141-1-Ig, Clone: 5A6B1) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 68141-1-Ig (TACSTD2 antibody) at dilution of 1:16000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).