For Research Use Only

USP36 Monoclonal antibody

Catalog Number:68165-1-lg Featured Product

1 Publications

57602

Q9P275

Full Name:

Calculated MW:

1123 aa, 123 kDa Observed MW: 150 kDa

ubiquitin specific peptidase 36



Basic Information

Applications

Catalog Number: 68165-1-lg

Size: 1000 µg/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG6426

Tested Applications: WB, ELISA

Cited Applications: WB

Species Specificity:

Human, Rat, Mouse **Cited Species:** human

GenBank Accession Number: **Purification Method:**

BC038983 Protein G purification GeneID (NCBI): CloneNo.:

1A1B3 **UNIPROT ID:** Recommended Dilutions:

WB 1:2000-1:10000

Positive Controls:

WB: U2OS cells, K-562 cells, HeLa cells, A549 cells, HEK-293 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells

Background Information

USP36 (Ubiquitin carboxyl-terminal hydrolase 36) is also named as KIAA1453, FLJ12851, DUB1 and belongs to the peptidase C19 family. It regulates the transcription and processing of rRNA by deubiquitylating various proteins involved in these processes, and the aberrant nucleolar structure in its absence is secondary to deficient ribosome biogenesis. USP36 plays a crucial role in regulating the structure and function of nucleoli (PMID:19208757). The ubiquitin-specific protease that is required in germline, epithelial, and intestinal stem cells (PMID:19039105). USP36 can be detected in 150 kDa (PMID:33852194)

Notable Publications

Author	Pubmed ID	Journal	Application
Gengchen Xie	38474064	Int J Mol Sci	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68165-1-1g (USP36 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68165-1-1g (USP36 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of USP36 antibody (68165-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP36 transfected HeLa cells.