

For Research Use Only

P5CS Monoclonal antibody

Catalog Number: 68184-1-Ig

Featured Product

1 Publications



Basic Information

Catalog Number:

68184-1-Ig

Concentration:

1000 ug/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG12100

GenBank Accession Number:

BC106054

GeneID (NCBI):

5832

UNIPROT ID:

P54886

Full Name:

aldehyde dehydrogenase 18 family, member A1

Calculated MW:

795 aa, 87 kDa

Observed MW:

87 kDa

Purification Method:

Protein A purification

CloneNo.:

3B1E11

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

IF

Species Specificity:

human, mouse, rat

Cited Species:

mouse

Positive Controls:

WB : U2OS cells, HAP1 cells, mouse colon tissue, rat colon tissue, A549 cells, MCF-7 cells, HeLa cells, LNCaP cells, HepG2 cells, K-562 cells

Background Information

Delta 1-pyrroline-5-carboxylate synthetase (P5CS), also named as GSAS, P5CS, PYCS, catalyzes the ATP and the NAD(P)H-dependent conversion of L-glutamate to glutamic gamma-semialdehyde (GSA) which is the metabolic precursor for proline biosynthesis(PMID:8761662). This protein includes 2 domains: glutamate 5-kinase and gamma-glutamyl phosphate reductase. In the N-terminal section, it belongs to the glutamate 5-kinase family and in the C-terminal section, it belongs to the gamma-glutamyl phosphate reductase family. Defects in ALDH18A1 are the cause of cutis laxa, autosomal recessive, type 3A (ARCL3A)(PMID:18478038). It has 2 isoforms produced by alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Keun Woo Ryu	39506109	Nature	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68184-1-Ig (P5CS antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



WB result of P5CS antibody (68184-1-Ig; 1:10000; room temperature for 1.5 hours) with wild-type and sg-P5CS transfected HAP1 cells.