For Research Use Only

P5CS Monoclonal antibody, PBS Only

Catalog Number:68184-1-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

3B1E11

Basic Information

Catalog Number: 68184-1-PBS Concentration:

1mg/ml Source: Mouse

lgG2b

Immunogen Catalog Number:

AG12100

795 aa, 87 kDa Observed MW: 87 kDa

member A1

Calculated MW:

BC106054

5832

P54886 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

aldehyde dehydrogenase 18 family,

Applications

Tested Applications:

WB, ELISA

Isotype:

Species Specificity:

human, mouse, rat

Background Information

Delta 1-pyrroline-5-carboxylate synthetase (P5CS), also named as GSAS, P5CS, PYCS, catalyzes the ATP and the $NAD(P) H-dependent\ conversion\ of\ L-glutamate\ to\ glutamic\ gamma-semial dehyde\ (GSA)\ which\ is\ the\ metabolic\ properties of\ the\ properties of\ the\ metabolic\ properties of\ the\ p$ precursor for proline biosynthesis (PMID:8761662). This protein includes 2 domains: glutamate 5-kinase and gammaglutamyl phosphate reductase. In the N-terminal section, it belongs to the glutamate 5-kinase family and in the Cterminal section, it belongs to the gamma-glutamyl phosphate reductase family. Defects in ALDH18A1 are the cause of cutis laxa, autosomal recessive, type 3A (ARCL3A)(PMID:18478038). It has 2 isoforms produced by alternative splicing.

Storage


Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only, pH7.3

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 68184-1-lg (P5CS antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 68184-1-PBS in a different storage buffer formulation.

WB result of P5CS antibody (68184-1-lg; 1:10000; room temperature for 1.5 hours) with wild-type and sg-P5CS transfected HAP1 cells. This data was developed using the same antibody clone with 68184-1-PBS in a different storage buffer formulation.